



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : C12N 15/30, 15/63, C07K 14/44, C12N 1/10, A61K 35/68, 38/17, C12P 21/08		A1	(11) International Publication Number: WO 95/06729
			(43) International Publication Date: 9 March 1995 (09.03.95)
(21) International Application Number: PCT/CA94/00482		(81) Designated States: AU, BR, CA, CN, FI, JP, KR, NO, NZ, RU, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 1 September 1994 (01.09.94)			
(30) Priority Data: 115,987 3 September 1993 (03.09.93) US		Published With international search report.	
(60) Parent Application or Grant (63) Related by Continuation US 08/115,987 (CIP) Filed on 3 September 1993 (03.09.93)			
(71) Applicant (for all designated States except US): MCGILL UNIVERSITY [CA/CA]; 845 Sherbrooke Street West, Montreal, Quebec H3A 2T5 (CA).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): MATLASHEWSKI, Gregory [CA/CA]; 2571 Chesnut Circle, St-Lazare, Quebec J0P 1V0 (CA); CHAREST, Hugues [CA/CA]; 1930 Sommet-Trinite, St-Bruno, Quebec H3V 4P6 (CA).			
(74) Agent: STEWART, Michael, I.; Sim & McBurney, Suite 701, 330 University Avenue, Toronto, Ontario M5G 1R7 (CA).			

(54) Title: DIFFERENTIALLY EXPRESSED *LEISHMANIA* GENES AND PROTEINS

[illegible]

**(57) Abstract**

Differentially expressed *Leishmania* genes and proteins are described. One differentially expressed gene (A2) is expressed at significantly elevated levels (more than about 10 fold higher) in the amastigote stage of the life cycle when the *Leishmania* organism is present in macrophages than in the free promastigote stage. The A2 gene encodes a 22 kD protein (A2 protein) that is recognized by kala-azar convalescent serum and has amino acid sequence homology with an S-antigen of *Plasmodium falciparum* Vietnamese isolate VI. Differentially expressed *Leishmania* genes and proteins have utility as vaccines, diagnostic reagents, as tools for the generation of immunological reagents and the generation of attenuated variants of *Leishmania*.

**BEST AVAILABLE COPY**

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

TITLE OF INVENTIONDIFFERENTIALLY EXPRESSED LEISHMANIA GENES AND PROTEINSFIELD OF INVENTION

5       The present invention is related to molecular cloning of Leishmania genes and, in particular, to the cloning of amastigote differentially expressed genes from Leishmania donovani.

BACKGROUND TO THE INVENTION

10       Leishmania protozoans are the causative agents of human leishmaniasis, which includes a spectrum of diseases ranging from self-healing skin ulcers to fatal visceral infections. Human leishmaniasis is caused by at least thirteen different species and subspecies of  
15 parasites of the genus Leishmania. Leishmaniasis has been reported from about eighty countries and probably some 400,000 new cases occur each year. Recently, the World Health Organization has reported 12 million people to be infected (ref. 1 - a listing of the references  
20 appears at the end of the disclosure).

L. donovani causes visceral leishmaniasis, also known as kala-azar. L. brasiliensis causes mucocutaneous leishmaniasis and L. major causes cutaneous leishmaniasis. Untreated visceral leishmaniasis is  
25 usually fatal and mucocutaneous leishmaniasis produces mutilation by destruction of the naso-oropharyngeal cavity and, in some cases, death.

      In addition, a major health problem has been created in areas of high infection when blood is collected for  
30 transfusion purposes. Since blood is a carrier of the parasites, blood from an infected individual may be unknowingly transferred to a healthy individual.

      The Leishmania protozoans exist as extracellular flagellated promastigotes in the alimentary tract of the  
35 sandfly in the free-living state, and are transmitted to the mammalian host through the bite of the insect vector. Once introduced, the promastigotes are taken up by

macrophages, rapidly differentiate into non-flagellated amastigotes and start to multiply within the phagolysosomal compartment. As the infected cells rupture, amastigotes subsequently infect other  
5 macrophages giving rise to the various symptoms associated with leishmaniasis (refs. 1 and 2). In this manner, it is the amastigote form of the parasite which is responsible for the pathology in humans.

While in the midgut of the insect, newly transformed  
10 promastigotes, functionally avirulent, progressively acquire capacity for infection and migrate to the mouthparts (ref. 3). This process, termed the metacyclogenesis, which occurs only in promastigotes, is concurrent with the differential expression of major  
15 surface glycoconjugates which mediate the migration of promastigotes in the alimentary tract and prepare the organism for the serum environment (refs. 4 and 5). In comparison, the promastigote to amastigote cytodifferentiation is a profound morphological and  
20 physiological transformation. During the promastigote to amastigote differentiation, the parasite loses its flagellum, rounds-up, changes its glycoconjugate coat (refs. 6, 7 and 8) and expresses a set of metabolic enzymes optimally active at low pH. The survival of the  
25 parasite inside the macrophage phagolysosome is associated with its ability to down-regulate many effector and accessory functions of its host cell, including oxygen metabolite-mediated killing and the capacity of the macrophage to act as an efficient antigen  
30 presenting cell (reviewed in, for example, ref. 9).

Leishmaniasis is, therefore, a serious disease and various types of vaccines against the disease have been developed, including live parasites; frozen promastigotes from culture; sonicated promastigotes; gamma-irradiated  
35 live promastigotes; and formalin-killed promastigotes treated with glucan (reviewed in, for example, ref. 10).

However, none of these approaches have provided satisfactory results.

The promastigote-amastigote differentiation is important to the establishment of infection. It would be desirable to identify genes and gene products that are differentially expressed when the amastigotes are present in macrophages.

Joshi, et al. describe L. donovani genes that are expressed at about two-fold higher in in vitro generated and maintained "amastigotes" compared to promastigotes (ref. 11).

#### SUMMARY OF THE INVENTION

The present invention is directed towards the provision of a Leishmania protein that is differentially expressed in the amastigote stage when the Leishmania organism is present within macrophages and genes encoding the differentially expressed protein. The amastigote differentially expressed gene and protein are useful for the preparation of vaccines against disease caused by Leishmania, the diagnosis of infection by Leishmania and as tools for the generation of immunological reagents and the generation of attenuated variants of Leishmania.

In accordance with one aspect of the present invention, there is provided a purified and isolated DNA molecule, the molecule comprising at least a portion coding for a differentially expressed gene of a Leishmania organism, the differentially expressed gene being expressed at an increased level when the amastigote form of the Leishmania organism is present within a macrophage. The increased level of expression maybe at least about a ten-fold increase in expression. In one embodiment of the present invention, the differentially expressed gene may be a virulence gene of the Leishmania organism and may be required for maintenance of infection by the amastigote form of the Leishmania organism.

In a further aspect of the invention, the differentially expressed virulence gene is functionally disabled by, for example, deletion or mutagenesis, such as insertional mutagenesis, to produce an attenuated Leishmania organism for use as, for example, a live vaccine. Conveniently, strains of Leishmania from which differentially expressed genes may be isolated include Leishmania donovani.

Further aspects of the invention include the protein encoded by the differentially expressed gene, and the use of the protein in vaccination and diagnosis. Additional aspects of the invention include an attenuated strain of Leishmania in which the virulence gene is disabled and a vaccine comprising the same.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a schematic outline of the amastigote cDNA library construction and differential screening with amastigote and promastigote-specific cDNA probes. An example of an amastigote-specific cDNA clone is indicated by an arrow on the colony hybridization autoradiogram;

Figure 2 shows a restriction enzyme and size analysis of Leishmania donovani amastigote-specific cDNA clones;

Figure 3 shows a Southern blot analysis of Leishmania donovani amastigote-specific cDNA clones;

Figure 4 shows a Northern blot analysis to demonstrate that A2-specific transcripts are present in amastigote-infected macrophages but not promastigotes;

Figure 5 shows a Southern blot analysis to demonstrate that A2 transcripts are encoded by a multigene family;

Figure 6 shows a restriction map of plasmid pGECO 90 that contains the L. donovani A2 gene;

Figure 7 shows a restriction map of a genomic clone of the A2 gene and its relationship to A2-related cDNAs;

Figure 8 shows the nucleotide sequence (SEQ ID NO: 1) and deduced amino acid sequence (SEQ ID NO: 2) of the open reading frame II (ORF II) of the Leishmania donovani A2 gene;

5        Figure 9 shows the homology between the Leishmania donovani A2 protein (SEQ ID NO: 2) and the Plasmodium falciparum S antigen (SEQ ID NO: 3) within the repeated subunits of these proteins;

10        Figure 10 shows the construction of a plasmid pET 16b/ORF II\* for expression of the A2 protein;

Figure 11 shows the presence of antibodies against A2 fusion protein in kala-azar immune serum by immunoprecipitation;

15        Figure 12 shows the specific recognition of A2 fusion protein by kala-azar sera by Western blot analysis; and

Figure 13 shows the results of Southern blot analysis of the separated chromosomes of different species and subspecies of Leishmania.

20        GENERAL DESCRIPTION OF THE INVENTION

Referring to Figure 1, there is illustrated a method used for isolating a Leishmania gene differentially expressed during the amastigote stage in the life cycle thereof. The method comprises the steps of (a)  
25        constructing a cDNA library from the Leishmania organism in the amastigote stage in the life cycle thereof; (b) constructing a first mixture of cDNA probes specific for the amastigote stage in the life cycle; (c) constructing a second mixture of cDNA probes specific for the  
30        promastigote stage in the life cycle; (d) separately probing the cDNA library with the amastigote and promastigote-specific cDNA probes in order to identify cDNA clones that are recognized by the amastigote mixture of cDNA probes but not the promastigote mixture of cDNA  
35        probes; and (e) isolating the cDNA clones identified in step (d).

The amastigote-specific cDNA clones identified by the above procedure can be further characterized by restriction enzyme analysis and their relatedness determined by Southern hybridization studies. To determine if cDNA clones identified by the above procedure represent amastigote-specific clones that are expressed at a higher level (more than about ten-fold higher) when the amastigote form of the Leishmania organism is present within macrophages, macrophages were infected with amastigotes and differentially-expressed gene transcripts were detected by Northern blot analysis. In an embodiment of the present invention, the differentially expressed Leishmania gene is L. donovani gene that is expressed at an increased level when the amastigote form of the Leishmania organism is present within a macrophage. The intracellular environment of the macrophage has an acidic pH of, for example, about 4.5. The differentially expressed genes include those having sequences, such as the DNA sequence set out in Figure 8 (SEQ ID No: 1) or its complementary strand; and DNA sequences which hybridize under stringent conditions to such DNA sequences. Such differentially expressed gene sequences include the A2 gene of L. donovani having the DNA sequence set out in Figure 8 and the invention includes a cDNA clone encoding the A2 gene depicted in Figure 8, which clone may be in the form of a plasmid, particularly that designated pGEC0 90 (Figure 6), which has ATCC accession number ATCC 75510.

The differentially expressed genes may encode proteins, such as the 22 kD A2 protein (SEQ ID No: 2), being encoded by the longest open reading frame (ORF II) of the A2 gene. Most of the predicted A2 protein is composed of a repetitive sequence consisting of a stretch of ten amino acids repeated nineteen times (Figure 8). Since each unit of this repeat contains two serines, two valines, two leucines and two prolines separated from



each other by five residues, the repeated region also may be considered as a stretch of five amino acids repeated thirty-eight times. The amino acid sequence of the A2 protein has homology with an S-antigen of Plasmodium falciparum (SEQ ID NO: 3), as shown in Figure 9. As with the L. donovani A2 protein, the carboxy-terminal portion of the S-antigen of P. falciparum Vietnamese isolate VI is composed of a stretch of eleven amino acids repeated nineteen times; the repeated units of both proteins are 50% identical and 80% homologous.

Life cycle stage specific genes from Leishmania may be isolated in the present invention. Some of these genes are required for transition between the life cycle stages and include virulence genes of the Leishmania parasite, such as virulence genes that are required for maintenance of infection by the amastigote form of the Leishmania organism. These virulence genes may be functionally disabled by, for example, deletion or mutation, including insertional mutagenesis and, furthermore, the wild-type Leishmania gene may be replaced by the functionally disabled gene. The virulence genes may be functionally disabled by, for example, replacing the A2 gene by a selectable antibiotic resistance gene by homologous recombination following transformation of the Leishmania organism with a fragment of DNA containing the antibiotic resistance gene flanked by 5'- and 3'- non-coding DNA sequences. This process can be used to generate attenuated variants of Leishmania and the residual pathogenicity of the attenuated variants can be assessed in mice and hamsters pigs. It is likely that deletion of genes that are selectively expressed in the human host environment (that being when the Leishmania organism is inside the macrophage cell) result in an attenuated strain which cannot survive in humans but generates a protective immune response. Attenuated strains of Leishmania would be useful as live vaccines

against the diseases caused by Leishmania and such attenuated strains form an aspect of the present invention.

Differentially expressed genes and proteins of  
5 Leishmania typified by the embodiments described herein are advantageous as:

- antigens for vaccination against the diseases caused by Leishmania.
- diagnostic reagents including hybridization  
10 probes, antigens and the means for producing specific antisera for (for example) detecting infection by Leishmania.
- target genes for functional  
15 disablement for the generation of attenuated Leishmania variants.

Vaccines comprising an effective amount of the protein encoded by the differentially expressed genes or of an attenuated strain of Leishmania and a physiologically-acceptable carrier therefor may utilize  
20 an adjuvant as the carrier and the protein may be presented to the immune system of the host in combination with an ISCOM or liposome. The vaccine may be formulated to be administered to a host in an injectable form, intranasally or orally, to immunize the host against  
25 disease.

#### BIOLOGICAL DEPOSITS

A plasmid pGECO 90 described and referred to herein was deposited with the American Type Culture Collection (ATCC) located at Rockville, Maryland, USA, pursuant to  
30 the Budapest Treaty on July 28, 1993 and prior to the filing of this application and assigned the ATCC accession number 75510. A diagram of this plasmid is shown in Figure 6. The plasmid contains the A2 gene of L. donovani described herein. The plasmid will become  
35 available to the public upon grant of a patent based upon this United States patent application. The invention

described and claimed herein is not to be limited in scope by the material deposited, since the deposited embodiment is intended only as an illustration of the invention. Any equivalent materials are within the scope  
5 of the invention.

#### EXAMPLES

The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific Examples. These  
10 Examples are described solely for purposes of illustration and are not intended to limit the scope of the invention. Changes in form and substitution of equivalents are contemplated as circumstances may suggest or render expedient. Although specific terms have been  
15 employed herein, such terms are intended in a descriptive sense and not for purposes of limitations.

Methods of molecular genetics and protein biochemistry used but not explicitly described in this disclosure and these Examples, are amply reported in the  
20 scientific literature and are well within the ability of those skilled in the art.

#### Example 1

This Example describes culturing and isolation of Leishmania organisms.

25 Amastigotes of the L. donovani Ethiopian LV9 strain were harvested from spleens of infected female gold Syrian hamsters and purified as described previously (ref. 12). Briefly, parasites were released from tissue using an homogenizer, the mixture was centrifuged three  
30 times at 100xg to remove cellular debris, and amastigotes were pelleted at 1500xg. The pellet was resuspended in 0.17 M sodium acetate to lyse contaminating red blood cells and amastigotes were recovered by centrifugation at 1500xg. Organisms were incubated at 37°C in complete  
35 RPMI medium (RPMI 1640 supplemented with 10% endotoxin free heat-inactivated FBS, 10 ml of 1M HEPES pH 7.3, 100

U of penicillin and 100 U of streptomycin per ml) for 18 hours prior to RNA extraction. After this period of incubation and multiple washes, the amastigote preparation was still physiologically active and relatively free of host cell contamination. To obtain promastigotes, LV9 strain amastigotes were allowed to differentiate in complete RPMI medium at 26°C, and cultured for at least seven days in the same medium before use (ref. 12).

Promastigotes of the L. donovani Sudanese strain 1S2D were cultivated and passaged in complete RPMI medium at 26°C. Amastigote-like organisms of the 1S2D strain were cultivated as described by Doyle et al. (ref. 13). The Sudanese strains 1S2D and 1S2D (wt) were obtained from Dr. S. Turco, the University of Kentucky, USA. The 1S2D (wt) promastigotes were adapted to grow in axenic conditions and had lost the ability to transform into infective promastigotes.

#### Example 2

This Example describes the preparation of and screening of a Leishmania cDNA library.

A method for isolating a Leishmania gene differentially expressed during the amastigote stage in the life cycle of the organism is illustrated in Figure 1.

Total RNA of amastigotes and promastigotes was prepared by the guanidinium isothiocyanate method using "RNAzol" (Trademark of Cinna/biotech Laboratories International Inc., Friendswood, TX); poly A<sup>+</sup> RNA was selected by oligo dT cellulose chromatography (grade 7:Pharmacia) as described by Sambrook et al. (ref. 14). A total of 10 µg of amastigote mRNA was used to construct an Eco RI/ Xho I unidirectional cDNA library of 10<sup>6</sup> clones in the "λ ZAP II" vector (Trademark of Stratagene); hemi-methylated cDNA was produced using the manufacturers reagents and protocols. About 40,000

amastigote and promastigote-specific clones of the primary library were screened differentially with amastigote and promastigote stage-specific gene probes. The cDNA probes were prepared using oligo dT<sub>12-18</sub> primer (Pharmacia) and M-MLV reverse transcriptase (BRL) following protocols previously described (ref. 15). Duplicate filters were hybridized with each probe for 18 h at 42°C in 50% formamide, 6X SSC, 5X Denhardt's solution, 5% dextran sulfate. Membranes then were washed twice at room temperature in 1X SSC for 20 min, twice at 55°C in 1X SSC, 0.1% SDS and then autoradiographed on "X-OMAT" films (Trademark of Kodak) with an intensifying screen for 18 to 72 hours. Such washing operation corresponds to stringent conditions of hybridization. Areas on the plates containing putative clones of interest were picked and the phage pools were submitted to a second round of screening. An example of an amastigote-specific cDNA clone is indicated by the arrow on the plaque hybridization autoradiogram of Figure 1.

Although cDNA clones representing promastigote-specific transcripts were more abundant than clones representing amastigote-specific transcripts, seven independent cDNA clones which only hybridized with amastigote-specific probes were isolated and termed 2, 3, 5, 6, 8, 9, 11. For each cDNA clone isolated, a Bluescript plasmid derivative was excised from the λZAP II recombinant phages in vivo using the helper phage R-408.

### Example 3

This Example describes the characterization of amastigote-specific cDNA clones.

The insert size of each of the Bluescript plasmids corresponding to the amastigote-specific cDNA clones was determined by restriction enzyme digestion and agarose gel electrophoresis (Figure 2). Recombinant plasmids (A2, A3, A5, A6, A8, A9 and A11) were digested with Eco

RI and Xho I to excise the cDNA inserts. Fragments were separated on a 1% agarose gel and stained with ethidium bromide. The cDNA inserts varied from 0.5 kb (A5) to 1.8 kb and A8 contained an internal Eco RI site. To  
5 determine if the amastigote-specific cDNA clones contain common sequences, Southern blot hybridization analysis of the Bluescript plasmids corresponding to the amastigote-specific cDNA clones was performed using clone A2 and clone A6 specific probes (Figure 3).

10 For Southern blot analysis, 10  $\mu$ g of total DNA was cut to completion with the restriction enzymes Eco RI and Xho I and separated on a 1% agarose gel. The restriction fragments were transferred to nylon membranes using standard procedures (ref. 16) and duplicates hybridized  
15 with  $\alpha$ -<sup>32</sup>P dCTP nick-translated probes representing the inserts of the cDNA clones A2 (0.9kb) or A6 (0.6kb). The A2 probe recognized five cDNAs (A2, A3, A8, A9 and A11) and the A6 cDNA only hybridized to itself. Thus, this Southern blot analysis indicated that cDNA clones A2, A3,  
20 A8, A9 and A11 contained homologous sequences but A5 and A6 were clones of unrelated amastigote-specific transcripts.

To confirm that the A2 series of clones represented Leishmania genes that were differentially expressed when  
25 the Leishmania organism is present in macrophages compared to expression in the free-living promastigotes, Northern blot analysis was performed. Total RNA was extracted from bone marrow-derived macrophages (BMM), L. donovani LV9-infected BMM (IBMM) and L. donovani LV9  
30 promastigotes (PRO). Murine bone marrow-derived macrophage cultures and L. donovani amastigote in vitro infections were carried out as previously described (ref. 12). The RNA species (15  $\mu$ g) were separated on an agarose gel and stained with ethidium bromide prior to  
35 transfer (Figure 4, right panel). The RNA was denatured by glyoxal treatment and transferred to a nylon membrane.

The Northern blot was hybridized with labelled cDNA A2 (0.9 kb) fragment, as previously described (ref. 12) (Figure 4, left panel). This probe recognized predominantly a 3.5 kb transcript present in amastigote-infected macrophages but not in promastigotes or in non-infected macrophages. This analysis showed that the A2 gene was differentially expressed at an increased level in amastigotes when they were present in macrophages compared to a free-living existence and that the increased expression was at least a ten fold increase.

#### Example 4

This Example describes the genomic arrangement and sequencing of the Leishmania donovani amastigote-specific A2 gene.

Regulation of transcription is one of the unusual features of the genetics of trypanosomatids. Copies of a gene or related genes are often clustered in tandem arrays on the same chromosome and a unique promoter region regulates expression of the cluster. Transcription leads to the synthesis of a polycistronic RNA molecule which is cleaved into monomeric units by trans-splicing prior to translation. The genomic arrangement of A2 related gene(s) was investigated by Southern blot analysis to determine whether it represents a multigene family. Total DNA was digested to completion with several restriction enzymes (E: Eco RI, S: Sal I, X: Xba I, C: Cla I, P: Pvu II). For double digests, the DNA was first cut to completion with Cla I or Pvu II, the DNA precipitated and resuspended in the appropriate buffer for the second digestion. Restriction fragments were separated on a 0.7% agarose gel, transferred to a nylon membrane and hybridized with a 0.5 kb Pst I/Xho I fragment of the A2 cDNA insert nick-translated with  $\alpha$ -<sup>32</sup>P dCTP. For each digest, the hybridization pattern displayed a series of bands of different intensities, clearly showing that many copies of the gene were present

in the genome (Figure 5). Moreover, common bands at about 6 to 8 kb for the Eco RI, Xba I and Sal I digests suggested an arrangement in tandem arrays. However, the presence of at least two other bands in each lane suggested that more than one cluster existed, each cluster being flanked by restriction fragments of different sizes. Alternatively, clusters also may carry copies of unrelated genes or intergenic regions of variable sizes.

5 To identify the protein coding region of A2, genomic clones carrying the A2 gene sequence were isolated. A partial genomic library containing 6 to 10 kb Eco RI fragments was constructed in the lambda ZAP II vector (Stratagene). More than 2,000 clones were screened on duplicate filters with probes prepared with the A2 cDNA using techniques and hybridization conditions described in Example 2. Eight clones were isolated and purified. Bluescript plasmid derivatives were excised from recombinant  $\lambda$  phages as for cDNA clones.

20 The 1.9 kb Xho I/ Eco RI insert fragment of the A2 Bluescript clone was subcloned into the Bluescript phagemids KS<sup>+</sup> and KS<sup>-</sup> for sequencing. Nested deletions were carried out on both plasmids using Exo III exonuclease and S1 nuclease. Sequencing reactions were performed on single-strand DNA templates using the M13K07 helper phage according to published procedures (ref. 17) with the Deaza G/A sequencing mixes (Pharmacia) and d<sup>35</sup>ATP or d<sup>35</sup>CTP radio-isotopes. Reactions were analysed on 6% denaturing gels. The inserts of the genomic clones were mapped with several restriction enzymes and displayed similar patterns, except some inserts were longer than others. One of these clones, pGECO 90 (as shown in Figure 6), was selected for further characterization. Figure 7 shows the restriction map of the insert of pGECO 90 and how it corresponds to the A2 related cDNAs. The restriction enzymes shown in Figure 7 are S: Sal I, P:



Pst I, O: Xho I, X: Xba I, E: Eco RI, M: Sma I. Plasmid pGECO 90 contained unique sites for Sal I and Xba I, but no Cla I site, and this was consistent with the Southern blot analysis shown in Figure 5. The DNA sequence  
5 flanking the Eco RI site on this genomic clone was determined and shown to correspond exactly to the related portion of the A8 cDNA, confirming that this fragment represented one unit of the tandem array.

The DNA sequence of the 1.9 kb Xho I/ Eco RI  
10 fragment of the pGECO 90 genomic clone corresponding to the 3.5 kb A2 transcript was determined (Figure 8) and compared to the cDNA's sequences. The longest open reading frame (ORF II) found was contained in the Xho I/  
Xba I 1.1 kb fragment and potentially encoded a 22 kD  
15 protein product (A2 protein). Stop codons were observed in two other frames and upstream from the initiating ATG. Most of this predicted A2 protein was composed of a repetitive sequence consisting of a stretch of ten amino acids repeated nineteen times. Since each unit of this  
20 repeat contains two serines, two valines, two leucines and two prolines separated from each other by five residues, the repeated region could also be considered as a stretch of five amino acids repeated thirty-eight times. The only hydrophobic domain was located at the  
25 amino terminal portion and may correspond to a signal peptide. The predicted amino acid sequence was compared with proteins reported in the Swiss-Prot database version using a Fasta algorithm (Canada Institute for Scientific and Technical Information: Scientific Numeric Database  
30 Service). The most striking identity was observed with an S-antigen of Plasmodium falciparum Vietnamese isolate VI. The alignment of the A2 protein sequence (A2) with the amino-terminal portion of the S-antigen of P. falciparum isolate VI is shown in Figure 9. Identical  
35 residues are indicated by dashes and homologous amino acids by dots. As with the L. donovani A2 protein, the

carboxy-terminal portion of this antigen of P. falciparum Vietnamese isolate IV is composed of a stretch of eleven amino acids repeated nineteen times. The repeated units of both proteins are 50% identical and 80% homologous.

5 The S-antigen, as the CS-antigens of Plasmodium, are proteins which are stage-specific, being expressed in the mammalian host but not in the insect host. Therefore, the A2 and S-antigen genes from unrelated human infectious protozoa are expressed specifically in the  
10 mammalian host and encode similar proteins. Thus, the A2 and S-antigen proteins may perform similar functions and may be required to enable these protozoa to survive in humans and functional disablement of the A2 sequences in L. donovani may be expected to result in a non-infective  
15 promastigote useful as a live attenuated vaccine for leishmaniasis.

#### Example 5

This Example describes the functional disablement of differentially expressed genes in Leishmania.

20 One approach for the development of attenuated strains of Leishmania is to functionally disable amastigote-specific genes (such as the A2 gene) from the Leishmania genome (by for example deletion) using homologous recombination. Deletion of genes from  
25 protozoa (such as Leishmania) has been described (ref. 18). This procedure involves cloning DNA fragments 5'- and 3'- to the A2 gene and constructing a plasmid vector that contains these flanking DNA sequences sandwiching a neomycin resistance gene. This 5'- neo 3'- fragment of  
30 DNA then is used to transform L. donovani promastigotes to G418 resistance. L. donovani is diploid and deletion one allele of the A2 gene in such G418 resistant strains can be determined by Southern blot hybridization using A2 specific probes. The second A2 allele then can be  
35 deleted by constructing a second deleting vector containing the 5'- and 3'- A2 flanking sequences

sandwiching a hygromycin resistance gene. Following transformation colonies are selected on medium containing G418 and hygromycin. Deletion of both copies of the A2 gene can be confirmed by Southern blot hybridization.

5 Example 6

This Example describes the expression of the L. donovani amastigote-specific A2 gene and the recognition of the A2 gene product by kala-azar immune sera.

To produce the A2 protein in a heterologous system, the coding region from the initiating ATG to the Xba I restriction site (see Figure 8) was subcloned in the pET 16B expression vector in frame with the HIS-TAG (Figure 10). The A2 fusion protein of 27 kD was produced in an in vitro transcription-translation assay (TNT system, Promega) using the pET16b/ORF II plasmid and a negative control pBluescript/p53 plasmid, encoding the human p53 protein. The in vitro translated HIS-TAG/A2 <sup>35</sup>S-labelled protein was immunoprecipitated with kala-azar immune serum and analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (Figure 11). Kala-azar is a term used to describe the disease caused by L. donovani. The kala-azar immune serum was obtained from a patient suffering from visceral leishmaniasis and reacted strongly against L. donovani antigens on ELISA.

25 In Figure 11, Lanes 1 and 2 contained the labelled proteins A2 and p53, respectively, prior to immunoprecipitation analysis. Lanes 3 and 4 contained proteins A2 and p53, respectively, immunoprecipitated with the kala-azar immune serum (L1) and Lanes 5 and 6 contained proteins A2 and p53, respectively, immunoprecipitated with a control human serum (TXC). The kala-azar serum did not react against the negative control protein human p53 but did immunoprecipitate the A2 gene-product. Neither of the proteins were immunoprecipitated by the control human serum. This analysis showed that the product of the L. donovani A2

gene was specifically recognized by kala-azar immune serum.

To confirm the specificity of the immune reaction, the pET 16b/ORF II plasmid coding for the recombinant A2 fusion protein and a negative control plasmid pET 16b with no insert, were introduced into E. coli. Expression was induced with IPTG, and total lysates of the recombinant E. coli cells separated by SDS-PAGE and analyzed by Western blot analysis using the kala-azar immune serum described above (see Figure 12). In Figure 12, Lane 1 contained E. coli/pET 16b cells and Lane 2 contained E. coli/pET 16b/ORF II cells. The kala-azar serum reacted specifically with protein products of 27.5 and 25 kD in the lysates of cells containing the pET 16b/ORF II plasmid (Lane 2). The 25 kD protein probably corresponded to the A2 protein without the HIS-TAG since the A2 sequence did contain its own initiating ATG. The serum did not react specifically with protein from E. coli lysates containing the control pET 16b plasmid (Lane 1). These data confirmed that the ORF II of the A2 gene encoded a L. donovani protein (A2) that was antigenic in patients with visceral leishmaniasis.

#### Example 7

This Example describes the Southern blot analysis of the isolated chromosomes of different species and subspecies of Leishmania.

Leishmania strains were obtained from American type Culture Collection, Rockville, Maryland, identified by their accession numbers as follows:

30		<u>ATCC</u>
	SF-2211: <u>L. donovani donovani</u> , strain MHOM/IN/80/DD8	50212
	SF-1881: <u>L. donovani infantum</u>	50134
	SF-1880: <u>L. donovani chagasi</u>	50133
35	SF-1882: <u>L. Braziliensis Braziliensis</u>	50135
	SF-1913: <u>L. Braziliensis panamensis</u>	50158
	SF-1871: <u>L. Braziliensis Guyanensis</u>	50126

	SF-1878: <u>L. Mexicana amazonensis</u>	50131
	SF-1911: <u>L. Mexicana mexicana</u>	50156
	SF-1864: <u>L. Major</u>	50122
	SF-1876: <u>L. tropica</u>	50129
5	SF-1861: <u>L. aetiopica</u>	50119

Sample blocks were initially prepared from the Leishmania strains, by the following protocol:

- 1) Promastigote cells were washed once in Hepes-NaCl buffer (21 mM HEPES pH 7.5, 137 mM NaCl, 5 mM KCl, 0.7 mM Na<sub>2</sub>PO<sub>4</sub> and 6 mM glucose) and resuspended at a density of  $5 \times 10^8$  in the same buffer.
- 2) Cells were diluted with 1 vol. of 1% low melting point agarose and 100 ul samples were allowed to cool down in sample holders at 4°C.
- 3) Blocks were transferred into lysis buffer (0.5 M EDTA pH 9.5, 1% sodium lauryl sarcosyl and 2 mg/ml of proteinase K) and incubated at 50°C for 18 h.
- 4) Blocks were kept at 4°C in 0.5 M EDTA.

Chromosomes were then separated from the sample blocks by Transverse Alternating Field Electrophoresis (TAFE) using a Geneline II System (Beckman instruments) under the following conditions:

- 1% agarose gel were prepared in 1X TAFE buffer (20X TAFE buffer consists in 0.45 M Tris-borate and 0.01 M EDTA). Electrophoresis were carried out at 350 mA for 36 h at 15°C.
- Electrophoresis conditions were:
  - Stage 1: 12 h, 40 s pulse time
  - Stage 2: 12 h, 100 s pulse time
  - Stage 3: 12 h, 160 s pulse time

Southern blots were then prepared from the chromosomal DNA by the following protocol:

- 1) Gels was soaked in 0.25 M HCl for 15 min. for a partial depurination of DNA.

2) DNA was denatured by an alkaline treatment (gels were soaked 0.5 N NaOH, 1.5 M NaCl for 45 min. with gentle shaking).

5 3) Gels were neutralized by soaking in 0.5 M Tris-Cl pH 7.0, 3 M NaCl for 45 min.

4) DNA was transferred to nylon membrane using a Vacugene XL (Pharmacia-LKB) for 2 h at 60 mbar in 10X SSC (1X SSC consists in 0.15 M NaCl, 0.015 M sodium citrate).

10 Hybridization next was carried out, as follows:

- Nylon membranes were prehybridized in 1 M NaCl, 1% sodium dodecyl sulfate (SDS), 10% dextran sulfate for 18 h. at 65°C for 2 h.

15 - Denatured probes were directly added to the hybridization buffer and blots were incubated for 18 h at 65°C.

- Membranes were washed twice in 2X SSC, 0.1% SDS at room temperature for 20 min. and twice in 0.5X SSC, 0.1% SDS for 20 min.

20 - Membranes were exposed on Kodak X-OMAT films with intensifying screens for 18 h.

The DNA probe used consisted for a pET16b/ORF II 1.1 kb Bam HI fragment agarose gel purified and labelled to high specificity with <sup>32</sup>P-dCTP (ICN;3000 ci/mMol) by nick-translation. This fragment contained the complete A2 protein coding region of the L. donovani A2 gene. The Southern blots obtained are shown in Figure 13.

30 The data provided by Figure 13 shows that the L. donovani A2 gene is present in all three species of L. donovani tested and two subspecies of L. mexicana. However, the A2 coding sequence was not found in L. tropica, L. major, L. braziliensis or L. aetiopica. From these results, it is apparent that the L. donovani A2 gene DNA is useful as a probe to detect specifically L. donovani and L. mexicana among Leishmania species. The L. donovani and L. mexicana species are usually

encountered at widely-different geographical locations, so the probe is specific for infection by the species present in a specific geographical location.

SUMMARY OF THE DISCLOSURE

5           In summary of this disclosure, the present invention provides differentially expressed genes and proteins of Leishmania, including the A2 gene expressed at significantly higher levels in the amastigote stage of the life cycle when the Leishmania organism is present in  
10   macrophages than in the promastigote stage. Modifications are possible within the scope of this invention.

REFERENCES

1. WHO, Tropical Disease Report, 1989. pp 85-92.
2. Turco, S.J., and Descoteaux, A. 1992. The Lipophosphoglycan of Leishmania parasites. Annu. Rev. Microbiol. 46:65-94.
3. Sacks, D.L. 1989. Metacyclogenesis in Leishmania promastigotes. Exp. Parasitology. 69:100-103.
4. Sacks D.L., and da Silva, R.P. 1987. The generation of infective stage L. major promastigotes is associated with the cell-surface expression and release of a developmentally regulated glycolipid. J. Immunol. 139:3099-3106.
5. Sacks, D.L., Brodin T.N., Turco, S.J. 1990. Developmental modification of the lipophosphoglycan from L. Major promastigotes during metacyclogenesis. Mol. Biochemical Parasitol. 42:225-234.
6. Medina-Acosta, E., Karess, R.E., Schwartz H., and Russell, D.G. 1989. The promastigote surface protease (gp63) of Leishmania is expressed but differentially processed and localized in the amastigote stage. Mol. Biochemical Parasitol. 37:263-274.
7. Turco, S.J. and Sacks, D.L. 1991. Expression of stage-specific lipophosphoglycan in Leishmania major amastigotes. Mol. Biochemical Parasitol. 45:91-100.
8. McConville, M.J., and Blackwell J.M. 1991. Developmental changes in the glycosylated phosphatidylinositols of L. donovani J. Biol. Chem. 260:15170-15179.
9. Bogdan, C., Röllinghoff M., and Solbach, W. 1990. Evasion strategies of Leishmania parasites. Parasitol. Today. 6:183-187.



10. Modabber, F. 1989. Experiences with vaccines against cutaneous leishmaniasis: of men and mice. *Parasitol.* 98:S49-S60.
11. Joshi, M., Dwyer, D.M., and Nakhasi, H.L. 1993. Cloning and characterization of differentially expressed genes from in vitro-grown "amastigotes" of Leishmania donovani. *Mol. Biochemical Parasitol.* 58:345-354.
12. Descoteaux, A., and Matlashewski, G. 1989. c-fos and tumor necrosis factor gene expression in Leishmania donovani-infected macrophages. *Mol.Cell. Biol.* 9:5223-5227.
13. Doyle, P.S. Engel, J.C., Pimenta, P.F.P. da Silva, P. and Dwyer. 1991. Leishmania donovani: Long-term culture of axenic amastigotes at 37°C. *Exp. Parasitol.* 73:326-334.
14. Sambrook, J., Fritsch, E.F., and Maniatis. 1989. Molecular cloning. A laboratory guide. Cold Spring Harbor Laboratories Press, New York. pp 7.26-7.29.
15. Sambrook, J., Fritsch, E.F., and Maniatis. 1989. Molecular cloning. A laboratory guide. Cold Spring Harbor Laboratories Press, New York. pp 10.44-10.45.
16. Sambrook, J., Fritsch, E.F., and Maniatis. 1989. Molecular cloning. A laboratory guide. Cold Spring Harbor Laboratories Press, New York. pp 9.38-9.40.
17. Sambrook, J., Fritsch, E.F., and Maniatis. 1989. Molecular cloning. A laboratory guide. Cold Spring Harbor Laboratories Press, New York. pp 4.48.
18. Cruz, A., and Beverley, S.M. 1990. Gene-replacement in parasitic protozoa. *Nature* 348:171-173.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

(A) NAME: McGill University  
(B) STREET: 845 Sherbrooke Street West  
(C) CITY: Montreal  
(D) STATE: Quebec  
(E) COUNTRY: Canada  
(F) POSTAL CODE (ZIP): H3A 2T5

(A) NAME: Gregory Matlashewski  
(B) STREET: 2571 Chestnut Circle  
(C) CITY: St-Lazare  
(D) STATE: Quebec  
(E) COUNTRY: Canada  
(F) POSTAL CODE (ZIP): J0P 1V0

(A) NAME: Hugues Charest  
(B) STREET: 1930 Sommet-Trinite  
(C) CITY: St-Bruno  
(D) STATE: Quebec  
(E) COUNTRY: Canada  
(F) POSTAL CODE (ZIP): H3V 4P6

(ii) TITLE OF INVENTION: DIFFERENTIALLY EXPRESSED LEISHMANIA GENES AND PROTEINS

(iii) NUMBER OF SEQUENCES: 4

## (iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1091 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GAGCTCCCCC AGCGACCCTC TCGGCAACGC GAGCGCCCCA GTCCCCCCAC GCACAAC TTT	60
GACCGAGCAC AATGAAGATC CGCAGCGTGC GTCCGCTTGT GGTGTTGCTG GTGTGCGTCG	120
CGGCGGTGCT CGCACTCAGC GCCTCCGCTG AGCCGCACAA GCGGCGCGTT GACGTCGGCC	180
CGCTCTCCGT TGGCCCGCAG TCCGTCGGCC CGCTCTCTGT TGGCCCGCAG GCTGTTGGCC	240
CGCTCTCCGT TGGCCCGCAG TCCGTCGGCC CGCTCTCTGT TGGCCCGCAG GCTGTTGGCC	300
CGCTCTCTGT TGGCCCGCAG TCCGTTGGCC CGCTCTCCGT TGGCCCGCTC TCCGTTGGCC	360
CGCAGTCTGT TGGCCCGCTC TCCGTTGGCT CGCAGTCCGT CGGCCCGCTC TCTGTTGGTC	420

CGCAGTCCGT CGGCCCCGCTC TCCGTTGGCC CGCAGGCTGT TGGCCCCGCTC TCCGTTGGCC	480
CGCAGTCCGT CGGCCCCGCTC TCTGTTGGCC CGCAGGCTGT TGGCCCCGCTC TCTGTTGGCC	540
CGCAGTCCGT TGGCCCCGCTC TCCGTTGGCC CGCAGTCTGT TGGCCCCGCTC TCCGTTGGCT	600
CGCAGTCCGT CGGCCCCGCTC TCTGTTGGTC CGCAGTCCGT CGGCCCCGCTC TCCGTTGGCC	660
CGCAGTCTGT CGGCCCCGCTC TCCGTTGGCC CGCAGTCCGT CGGCCCCGCTC TCCGTTGGTC	720
CGCAGTCCGT TGGCCCCGCTC TCCGTTGGCC CGCAGTCCGT TGACGTTTCT CCGGTGTCTT	780
AAGGCTCGGC GTCCGCTTTC CCGGTGTGCGT AAAGTATATG CCATGAGGCA TGGTGACGAG	840
GCAAACCTTG TCAGCAATGT GGCATTATCG TACCCGTGCA AGAGCAACAG CAGAGCTGAG	900
TGTTCAAGTG GCCACAGCAC CACGCTCCTG TGACACTCCG TGGGGTGTGT GTGACCTTGG	960
CTGCTGTTGC CAGGCGGATG AACTGCGAGG GCCACAGCAG CGCAAGTGCC GCTTCCAACC	1020
TTGCGACTTT CACGCCACAG ACGCATAGCA GCGCCCTGCC TGTCGCGGCG CATGCGGGCA	1080
AGCCATCTAG A	1091

## (2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 711 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

ATGAAGATCC GCAGCGTGCG TCCGCTTGTC GTGTTGCTGG TGTGCGTCGC GCGGGTGCTC	60
GCACTCAGCG CCTCCGCTGA GCCGCACAAG GCGGCCGTTG ACGTCGGCCC GCTCTCCGTT	120
GGCCCGCAGT CCGTCGGCCC GCTCTCTGTT GGCCCGCAGG CTGTTGGCCC GCTCTCCGTT	180
GGCCCGCAGT CCGTCGGCCC GCTCTCTGTT GGCCCGCAGG CTGTTGGCCC GCTCTCTGTT	240
GGCCCGCAGT CCGTTGGCCC GCTCTCCGTT GGCCCGCTCT CCGTTGGCCC GCAGTCTGTT	300
GGCCCGCTCT CCGTTGGCTC GCAGTCCGTC GGCCCGCTCT CTGTTGGTCC GCAGTCCGTC	360
GGCCCGCTCT CCGTTGGCCC GCAGGCTGTT GGCCCGCTCT CCGTTGGCCC GCAGTCCGTC	420
GGCCCGCTCT CTGTTGGCCC GCAGGCTGTT GGCCCGCTCT CTGTTGGCCC GCAGTCCGTT	480
GGCCCGCTCT CCGTTGGCCC GCAGTCTGTT GGCCCGCTCT CCGTTGGCTC GCAGTCCGTC	540
GGCCCGCTCT CTGTTGGTCC GCAGTCCGTC GGCCCGCTCT CCGTTGGCCC GCAGTCTGTC	600
GGCCCGCTCT CCGTTGGCCC GCAGTCCGTC GGCCCGCTCT CCGTTGGTCC GCAGTCCGTT	660
GGCCCGCTCT CCGTTGGCCC GCAGTCCGTT GACGTTTCTC CCGTGTCTTA A	711

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 236 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```

Met Lys Ile Arg Ser Val Arg Pro Leu Val Val Leu Leu Val Cys Val
 1           5           10           15
Ala Ala Val Leu Ala Leu Ser Ala Ser Ala Glu Pro His Lys Ala Ala
 20           25           30
Val Asp Val Gly Pro Leu Ser Val Gly Pro Gln Ser Val Gly Pro Leu
 35           40           45
Ser Val Gly Pro Gln Ala Val Gly Pro Leu Ser Val Gly Pro Gln Ser
 50           55           60
Val Gly Pro Leu Ser Val Gly Pro Gln Ala Val Gly Pro Leu Ser Val
 65           70           75           80
Gly Pro Gln Ser Val Gly Pro Leu Ser Val Gly Pro Leu Ser Val Gly
 85           90           95
Pro Gln Ser Val Gly Pro Leu Ser Val Gly Ser Gln Ser Val Gly Pro
100           105           110
Leu Ser Val Gly Pro Gln Ser Val Gly Pro Leu Ser Val Gly Pro Gln
115           120           125
Ala Val Gly Pro Leu Ser Val Gly Pro Gln Ser Val Gly Pro Leu Ser
130           135           140
Val Gly Pro Gln Ala Val Gly Pro Leu Ser Val Gly Pro Gln Ser Val
145           150           155           160
Gly Pro Leu Ser Val Gly Pro Gln Ser Val Gly Pro Leu Ser Val Gly
165           170           175
Ser Gln Ser Val Gly Pro Leu Ser Val Gly Pro Gln Ser Val Gly Pro
180           185           190
Leu Ser Val Gly Pro Gln Ser Val Gly Pro Leu Ser Val Gly Pro Gln
195           200           205
Ser Val Gly Pro Leu Ser Val Gly Pro Gln Ser Val Gly Pro Leu Ser
210           215           220
Val Gly Pro Gln Ser Val Asp Val Ser Pro Val Ser
225           230           235

```

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 269 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Pro Gly Ser Glu Gly Pro Lys Gly Thr Gly Gly Pro Gly Ser Glu Gly  
 1 5 10 15  
 Pro Lys Gly Thr Gly Gly Pro Gly Ser Glu Gly Pro Lys Gly Thr Gly  
 20 25 30  
 Gly Pro Gly Ser Glu Gly Pro Lys Gly Thr Gly Gly Pro Gly Ser Glu  
 35 40 45  
 Gly Pro Lys Gly Thr Gly Gly Pro Gly Ser Glu Gly Pro Lys Gly Thr  
 50 55 60  
 Gly Gly Pro Gly Ser Glu Gly Pro Lys Gly Thr Gly Gly Pro Gly Ser  
 65 70 75 80  
 Glu Gly Pro Lys Gly Thr Gly Gly Pro Gly Ser Glu Gly Pro Lys Gly  
 85 90 95  
 Thr Gly Gly Pro Gly Ser Glu Gly Pro Lys Gly Thr Gly Gly Pro Gly  
 100 105 110  
 Ser Glu Gly Pro Lys Gly Thr Gly Gly Pro Gly Ser Glu Gly Pro Lys  
 115 120 125  
 Gly Thr Gly Gly Pro Gly Ser Glu Gly Pro Lys Gly Thr Gly Gly Pro  
 130 135 140  
 Gly Ser Glu Gly Pro Lys Gly Thr Gly Gly Pro Gly Ser Glu Gly Pro  
 145 150 155 160  
 Lys Gly Thr Gly Gly Pro Gly Ser Glu Gly Pro Lys Gly Thr Gly Gly  
 165 170 175  
 Pro Gly Ser Glu Ser Pro Lys Gly Thr Gly Gly Pro Gly Ser Glu Gly  
 180 185 190  
 Pro Lys Gly Thr Gly Gly Pro Gly Ser Glu Gly Pro Lys Gly Thr Gly  
 195 200 205  
 Pro Lys Gly Thr Gly Gly Pro Gly Ser Glu Ala Gly Thr Glu Gly Pro  
 210 215 220  
 Lys Gly Thr Gly Gly Pro Gly Ser Glu Ala Gly Thr Glu Gly Pro Lys  
 225 230 235 240  
 Gly Thr Gly Gly Pro Gly Ser Gly Gly Glu His Ser His Asn Lys Lys  
 245 250 255  
 Lys Ser Lys Lys Ser Ile Met Asn Met Leu Ile Gly Val  
 260 265

CLAIMS

What we claim is:

1. An isolated and purified chromosomal DNA molecule, the molecule comprising at least a portion coding for a differentially expressed gene of a Leishmania organism, the differentially expressed gene being expressed at an increased level when the amastigote form of the Leishmania organism is present within a macrophage in comparison to the promastigote form of the Leishmania organism.
2. The DNA molecule of claim 1 wherein the increased level is at least about a ten fold increase.
3. The DNA molecule of claim 1 wherein the differentially expressed gene is a virulence gene of the Leishmania organism.
4. The DNA molecule of claim 3 wherein the virulence gene is required for maintenance of infection by the amastigote form of the Leishmania organism.
5. The DNA molecule of claim 1 wherein the differentially expressed gene encodes a protein.
6. The DNA molecule of claim 1 wherein the Leishmania organism is Leishmania donovani.
7. The DNA molecule of claim 1 wherein the differentially expressed gene has the DNA coding sequence set out in Figure 8 or its complementary strand or a DNA molecule coding for a differentially expressed gene of a Leishmania organism which hybridizes under stringent conditions thereto.
8. An isolated and purified DNA fragment having the nucleotide sequence:

```

GAGCTCCCC AGCGACCCTC TCGGCAACGC GAGCGCCCCA GTCCCCCCAC GCACAACTTT   60
GACCGAGCAC AATGAAGATC CGCAGCGTGC GTCCGCTTGT GGTGTTGCTG GTGTGCGTCG   120
CGGCGGTGCT CGCACTCAGC GCCTCCGCTG AGCCGCACAA GGCGGCCGTT GACGTGGGCC   180
CGCTCTCCGT TGGCCCGCAG TCCGTGCGCC CGCTCTCTGT TGGCCCGCAG GCTGTTGGCC   240
CGCTCTCCGT TGGCCCGCAG TCCGTGCGCC CGCTCTCTGT TGGCCCGCAG GCTGTTGGCC   300
CGCTCTCTGT TGGCCCGCAG TCCGTTGGCC CGCTCTCCGT TGGCCCGCTC TCCGTTGGCC   360
CGCAGTCTGT TGGCCCGCTC TCCGTTGGCT CGCAGTCCGT CGGCCCGCTC TCTGTTGGTC   420
CGCAGTCCGT CGGCCCGCTC TCCGTTGGCC CGCAGGCTGT TGGCCCGCTC TCCGTTGGCC   480
CGCAGTCCGT CGGCCCGCTC TCTGTTGGCC CGCAGGCTGT TGGCCCGCTC TCTGTTGGCC   540

```

29

CGCAGTCCGT TGGCCCGCTC TCCGTTGGCC CGCAGTCTGT TGGCCCGCTC TCCGTTGGCT 600  
 CGCAGTCCGT CGGCCCGCTC TCTGTTGGTC CGCAGTCCGT CGGCCCGCTC TCCGTTGGCC 660  
 CGCAGTCTGT CGGCCCGCTC TCCGTTGGCC CGCAGTCCGT CGGCCCGCTC TCCGTTGGTC 720  
 CGCAGTCCGT TGGCCCGCTC TCCGTTGGCC CGCAGTCCGT TGACGTTTCT CCGGTGTCTT 780  
 AAGGCTCGGC GTCCGCTTTC CGGTGTGCGT AAAGTATATG CCATGAGGCA TGGTGACGAG 840  
 GCAAACCTTG TCAGCAATGT GGCATTATCG TACCCGTGCA AGAGCAACAG CAGAGCTGAG 900  
 TGTTACAGTG GCCACAGCAC CACGCTCCTG TGACACTCCG TGGGGTGTGT GTGACCTTGG 960  
 CTGCTGTTGC CAGGCGGATG AACTGCGAGG GCCACAGCAG CGCAAGTGCC GCTTCCAACC 1020  
 TTGCGACTTT CACGCCACAG ACGCATAGCA GCGCCCTGCC TGTCGCGGCG CATGCGGGCA 1080  
 AGCCATCTAG A 1091

(SEQ ID NO: 1), or its complementary strand, or a DNA molecule coding for a differentially-expressed gene of a Leishmania organism which hybridizes thereto under stringent conditions.

9. An isolated and purified DNA fragment having the nucleotide sequence:

ATGAAGATCC GCAGCGTGC TCCGCTTGTG GTGTTGCTGG TGTGCGTCGC GGCGGTGCTC 60  
 GCACTCAGCG CCTCCGCTGA GCCGCACAA GCGGCCGTTG ACGTCGGCCC GCTCTCCGTT 120  
 GGCCCGCAGT CCGTCGGCCC GCTCTCTGTT GGCCCGCAGG CTGTTGGCCC GCTCTCCGTT 180  
 GGCCCGCAGT CCGTCGGCCC GCTCTCTGTT GGCCCGCAGG CTGTTGGCCC GCTCTCTGTT 240  
 GGCCCGCAGT CCGTTGGCCC GCTCTCCGTT GGCCCGCTCT CCGTTGGCCC GCAGTCTGTT 300  
 GGCCCGCTCT CCGTTGGCTC GCAGTCCGTC GGCCCGCTCT CTGTTGGTCC GCAGTCCGTC 360  
 GGCCCGCTCT CCGTTGGCCC GCAGGCTGTT GGCCCGCTCT CCGTTGGCCC GCAGTCCGTC 420  
 GGCCCGCTCT CTGTTGGCCC GCAGGCTGTT GGCCCGCTCT CTGTTGGCCC GCAGTCCGTT 480  
 GGCCCGCTCT CCGTTGGCCC GCAGTCTGTT GGCCCGCTCT CCGTTGGCTC GCAGTCCGTC 540  
 GGCCCGCTCT CTGTTGGTCC GCAGTCCGTC GGCCCGCTCT CCGTTGGCCC GCAGTCTGTC 600  
 GGCCCGCTCT CCGTTGGCCC GCAGTCCGTC GGCCCGCTCT CCGTTGGTCC GCAGTCCGTT 660  
 GGCCCGCTCT CCGTTGGCCC GCAGTCCGTT GACGTTTCTC CCGTGTCTTA A 711

(SEQ ID NO: 2), or its complementary strand, or a DNA molecule coding for a differentially-expressed gene of a Leishmania organism which hybridizes thereto under stringent conditions.

10. An isolated and purified DNA fragment encoding the amino acids sequence:

Met	Lys	Ile	Arg	Ser	Val	Arg	Pro	Leu	Val	Val	Leu	Leu	Val	Cys	Val
1					5					10				15	
Ala	Ala	Val	Leu	Ala	Leu	Ser	Ala	Ser	Ala	Glu	Pro	His	Lys	Ala	Ala
					20					25				30	
Val	Asp	Val	Gly	Pro	Leu	Ser	Val	Gly	Pro	Gln	Ser	Val	Gly	Pro	Leu
					35					40				45	

```

Ser Val Gly Pro Gln Ala Val Gly Pro Leu Ser Val Gly Pro Gln Ser
 50                      55                      60
Val Gly Pro Leu Ser Val Gly Pro Gln Ala Val Gly Pro Leu Ser Val
65                      70                      75                      80
Gly Pro Gln Ser Val Gly Pro Leu Ser Val Gly Pro Leu Ser Val Gly
      85                      90                      95
Pro Gln Ser Val Gly Pro Leu Ser Val Gly Ser Gln Ser Val Gly Pro
      100                      105                      110
Leu Ser Val Gly Pro Gln Ser Val Gly Pro Leu Ser Val Gly Pro Gln
      115                      120                      125
Ala Val Gly Pro Leu Ser Val Gly Pro Gln Ser Val Gly Pro Leu Ser
      130                      135                      140
Val Gly Pro Gln Ala Val Gly Pro Leu Ser Val Gly Pro Gln Ser Val
145                      150                      155                      160
Gly Pro Leu Ser Val Gly Pro Gln Ser Val Gly Pro Leu Ser Val Gly
      165                      170                      175
Ser Gln Ser Val Gly Pro Leu Ser Val Gly Pro Gln Ser Val Gly Pro
      180                      185                      190
Leu Ser Val Gly Pro Gln Ser Val Gly Pro Leu Ser Val Gly Pro Gln
      195                      200                      205
Ser Val Gly Pro Leu Ser Val Gly Pro Gln Ser Val Gly Pro Leu Ser
      210                      215                      220
Val Gly Pro Gln Ser Val Asp Val Ser Pro Val Ser
225                      230                      235

```

(SEQ ID NO: 3), or its complementary strand, or a DNA molecule coding for a differentially-expressed gene of a Leishmania organism which hybridizes thereto under stringent conditions.

11. A recombinant plasmid adapted for transformation of a microbial host, the recombinant plasmid comprising a plasmid vector into which a DNA segment comprising the purified and isolated DNA molecule of any one of claims 1 to 10 has been inserted.

12. The recombinant plasmid of claim 11 which is plasmid pGECO 90 having ATCC accession number 75510.

13. A purified protein encoded by a differentially expressed gene of a Leishmania organism, the differentially expressed gene being expressed at an increased level when the amastigote form of the Leishmania organism is present within a macrophage.



14. The purified protein of claim 13 wherein the increased level is at least about a ten fold increase.
15. The protein of claim 13 wherein the differentially expressed gene is a virulence gene of the Leishmania organism.
16. The protein of claim 15 wherein the virulence gene is required for maintenance of infection by the amastigote form of the Leishmania organism.
17. The protein of claim 13 wherein the differentially expressed gene has the DNA sequence set out in Figure 8 or its complementary strand, or a DNA sequence coding for a differentially expressed gene of a Leishmania organism which hybridizes under stringent conditions thereto.
18. The protein of claim 13 wherein the Leishmania organism is Leishmania donovani.
19. An attenuated strain of Leishmania wherein the virulence gene has been functionally disabled.
20. The attenuated strain of claim 19 wherein the virulence gene has been functionally disabled by deletion.
21. The attenuated strain of claim 19 wherein the virulence gene has been functionally disabled by mutagenesis thereof.
22. The attenuated strain of claim 21 wherein the virulence gene has been functionally disabled by insertional mutagenesis.
23. The attenuated strain of claim 19 wherein the differentially expressed virulence gene has the DNA sequence set out in Figure 8 or its complementary strand, or a DNA sequence coding for a differentially expressed gene of a Leishmania organism which hybridizes under stringent conditions thereto.
24. A vaccine to provide protective immunity to a host against disease caused by a Leishmania organism, comprising an effective amount of the protein claimed in claim 13 and a physiologically-acceptable carrier therefor.

25. The vaccine of claim 24 wherein the carrier comprises an adjuvant.

26. The vaccine of claim 24 wherein the protein is presented to the immune system of the host in combination with an ISCOM or a liposome.


27. A live vaccine to provide protective immunity to a host against disease caused by a Leishmania organism, comprising an effective amount of the attenuated strain of Leishmania wherein the virulence gene has been functionally disabled and a physiologically-acceptable carrier therefor.

28. The vaccine of claim 24 or 27 formulated to be administered in an injectable form, intranasally or orally.

29. A method of immunizing a host against disease caused by a Leishmania organism, which comprises administering to the host an effective amount of vaccine claimed in any one of claims 24 or 27.

30. An antibody raised against the protein of claim 13.

1 / 12



*L. donovani* LV9  
infected cells

Extraction of amastigotes

incubation at 26°C  
(7 days)

incubation at 37°C  
(18 h.)



RNA extraction;  
poly A selection



-hemimethylated cDNA  
-unidirectional insertion  
in a Zap II vector

promastigote  
cDNA probe

amastigote  
cDNA probe




FIG.1.

SUBSTITUTE SHEET

2 / 12

2 3 5 6 8 9 11




**FIG.2.**

(probes)

$$\begin{array}{c} A2 \\ \hline 2 \ 3 \ 5 \ 6 \ 8 \ 9 \ 11 \\ \hline \end{array}$$

A6  
2 3 5 6 8 9 11



**FIG. 3.**




FIG.4.




FIG.5.

**FIG. 6.**

5/12




FIG.7

6/12

FIG.8A.

XHO I

GAGCTCCCCCAGCGACCCCTCTCGGCAACGGAGCGCCCCAGTCCCCCAGCACAACTTTGACCCGAGCACA

ORF II

Met Lys Ile Arg Ser Val Arg Pro Leu Val Leu Val Leu Val Cys Val Ala Val Leu Ala Leu  
 1, ATG AAG ATC CGC AGC GTG CGT CCG CTT GTG TCG TTT GTG TGC GCG GCG GTG CTC GCA CTC

Ser Ala Ser Ala Glu Pro His Lys Ala Ala Val Asp  
 67 AGC GCC TCC GCT GAG CCG CAC AAG GCG GCC GTT GAC

Val Gly Pro Leu Ser Val Gly Pro  
 103 GTC GGC CCG CTC TCC GTT GGC CCG

Gln Ser Val Gly Pro Leu Ser Val Gly Pro  
 127 CAG TCC GTC GGC CCG CTC TCT GTT GGC CCG

Gln Ala Val Gly Pro Leu Ser Val Gly Pro  
 157 CAG GCT GTT GGC CCG CTC TCC GTT GGC CCG

Gln Ser Val Gly Pro Leu Ser Val Gly Pro  
 187 CAG TCC GTC GGC CCG CTC TCT GTT GGC CCG

Gln Ala Val Gly Pro Leu Ser Val Gly Pro  
 217 CAG GCT GTT GGC CCG CTC TCC GTT GGC CCG

Gln Ser Val Gly Pro Leu Ser Val Gly Pro Leu Ser Val Gly Pro  
 247 CAG GCT GTT GGC CCG CTC TCC GTT GGC CCG CTC TCC GTT GGC CCG

SUBSTITUTE SHEET



7/12

FIG.8B.

Gln Ser Val Gly Pro Leu Ser Val Gly Ser  
292 CAG TCT GTC GGC CCG CTC TCC GTT GGC TCG

Gln Ser Val Gly Pro Leu Ser Val Gly Pro  
322 CAG TCC GTC GGC CCG CTC TCT GTT GGT CCG

Gln Ser Val Gly Pro Leu Ser Val Gly Pro  
352 CAG TCC GTC GGC CCG CTC TCT GTT GGC CCG

Gln Ala Val Gly Pro Leu Ser Val Gly Pro  
382 CAG GCT GTT GGC CCG CTC TCC GTT GGC CCG

Gln Ser Val Gly Pro Leu Ser Val Gly Pro  
412 CAG TCC GTC GGC CCG CTC TCT GTT GGC CCG

Gln Ala Val Gly Pro Leu Ser Val Gly Pro  
442 CAG GCT GTT GGC CCG CTC TCT GTT GGC CCG

Gln Ser Val Gly Pro Leu Ser Val Gly Pro  
472 CAG TCC GTT GGC CCG CTC TCC GTT GGC CCG

Gln Ser Val Gly Pro Leu Ser Val Gly Ser  
502 CAG TCT GTT GGC CCG CTC TCC GTT GGC TCG

Gln Ser Val Gly Pro Leu Ser Val Gly Pro  
532 CAG TCC GTC GGC CCG CTC TCT GTT GGT CCG

Gln Ser Val Gly Pro Leu Ser Val Gly Pro  
562 CAG TCC GTC GGC CCG CTC TCC GTT GGC CCG

SUBSTITUTE SHEET

8/12

FIG.8C.

Gln Ser Val Gly Pro Leu Ser Val Gly Pro  
 592 CAG TCT GTC GGC CCG CTC TCC GTT GGC CCG

Gln Ser Val Gly Pro Leu Ser Val Gly Pro  
 622 CAG TCC GTC GGC CCG CTC TCC GTT GGT CCG

Gln Ser Val Gly Pro Leu Ser Val Gly Pro  
 652 CAG TCC GTT GGC CCG CTC TCC GTT GGC CCG

Gln Ser Val  
 682 CAG TCC GTC

Asp Val Ser Pro Val Ser \*\*\*

691 GAC GTT TCT CCG GTG TCT TAAGGCTCGGCGTCCGCTTCCGGTGTGCGTAAAGTATATGCCATGAGGCATGGTGACGAGGCAAAC  
 776 CTTGTCAGCAATGTGGCATTATCGTACCCGTGCAAGAGCAACAGCAGAGCTGAGTGTTCAGGTGGCCACAGCACCCACGCTCCTGTGACACT  
 867 CCGTGGGGTGTGTGACCTTGGCTGTGTTGCCAGGGGGATGAACCTGCCAGGGGCCACAGCAGCGCAAGTGCCCGCTTCCAAACCTTGGGACT  
 958 TTCACGCCACAGACGATAGCAGCGCCCTGCTGTCGGGGCGCATGCGGGCAAGCCATCTAGA

XBA I

SUBSTITUTE SHEET

9/12

**FIG. 9.**

```

A2          10      20      30      40      50
      MKIRSVRPLVLLVCVAAVLALSASAEPHKAAVDVGPLSVGPQSV-GPLSVG
Sant_P      PGSEGPKGTTGGPGSEGPKGTTGGPGSEGPKGTTGGPGSEGPKGTTGGPGSEGP
100      120      130      140      150      160

A2          60      70      80      90      100
      PQAV-GPLSVGPQSV-GPLSVGPQAV-GPLSVGPQSVGPLSVGPQSV-GPLSVGS
|::: || ||::: || ||::: || ||::: || ||::: || ||::: || ||::: ||
Sant_P      PKGTGGPGSEGPKGTTGGPGSEGPKGTTGGPGSEGPKGTTGGPGSEGP
170      180      190      200      210      220

A2          60      70      80      90      100
      QSV-GPLSVGPQSV-GPLSVGPQAV-GPLSVGPQSV-GPLSVGPQAV-GPLSVGPQSV-G
::: || ||::: || ||::: || ||::: || ||::: || ||::: || ||::: ||
Sant_P      KGTGGPGSEGPKGTTGGPGSEGPKGTTGGPGSEGPKGTTGGPGSEGPKGTTGG
230      240      250      260      270      280

A2          60      70      80      90      100
      PLSVGPQSV-GPLSVGSQSV-GPLSVGPQSVGPLSVGSQSVGPLSVGSQSVGPLSVGSQS
| |::: || ||::: || ||::: || ||::: || ||::: || ||::: || |::: |
Sant_P      PGSESPKGTGGPGSEGPKGTTGGPGSEGPKGTTGGPGSEAGTEGPKGTGGPGSEAGT
290      300      310      320      330      340

A2          60      70      80      90      100
      VGPLSVGPQSVDVSPVS
| |::: |
Sant_P      EGPKGTTGGPGSGGEHSHNKKKKSSIMNMLIGV
350      360      370

```




FIG.10.




FIG.11.

SUBSTITUTE SHEET

11 / 12




FIG.12.

12 / 12




FIG.13.

## INTERNATIONAL SEARCH REPORT

 Internati Application No  
 PCT/CA 00482

 A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 6 C12N15/30 C12N15/63 C07K14/44 C12N1/10 A61K35/68  
 A61K38/17 C12P21/08

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

 Minimum documentation searched (classification system followed by classification symbols)  
 IPC 6 C07K A61K C12N C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MOLECULAR AND BIOCHEMICAL PARASITOLOGY, vol.58, no.2, 1993 pages 345 - 354 MANJU JOSHI ET AL. 'Cloning and characterization of differentially expressed genes from in vitro-grown 'amastigotes' of Leishmania donovani' cited in the application see abstract see page 348, left column, last paragraph - right column, paragraph 1 see page 349, left column, paragraph 2 - right column, paragraph 1 see page 351, left column, paragraph 2 - right column, paragraph 1 --- -/--	1,5,6, 13,15,18

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

## \* Special categories of cited documents:

 "A" document defining the general state of the art which is not  
 considered to be of particular relevance

 "E" earlier document but published on or after the international  
 filing date

 "L" document which may throw doubts on priority claim(s) or  
 which is cited to establish the publication date of another  
 citation or other special reason (as specified)

 "O" document referring to an oral disclosure, use, exhibition or  
 other means

 "P" document published prior to the international filing date but  
 later than the priority date claimed

 "T" later document published after the international filing date  
 or priority date and not in conflict with the application but  
 cited to understand the principle or theory underlying the  
 invention

 "X" document of particular relevance; the claimed invention  
 cannot be considered novel or cannot be considered to  
 involve an inventive step when the document is taken alone

 "Y" document of particular relevance; the claimed invention  
 cannot be considered to involve an inventive step when the  
 document is combined with one or more other such docu-  
 ments, such combination being obvious to a person skilled  
 in the art.

"&amp;" document member of the same patent family

Date of the actual completion of the international search

22 November 1994

Date of mailing of the international search report

14-12-1994

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax (+31-70) 340-3016

Authorized officer

Montero Lopez, B

## INTERNATIONAL SEARCH REPORT

 International Publication No  
 PCT/C/00482

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	THE JOURNAL OF PROTOZOOLOGY, vol.26, no.3, August 1979 page 49A ESTHER MARVA ET AL. 'Vaccination of mice with chemically attenuated Leishmani tropica' see abstract n. 140 ---	19
A	JOURNAL OF IMMUNOLOGY., vol.140, no.7, 1 April 1988, BALTIMORE US pages 2406 - 2414 ALFRED A. PAN ET AL. 'Monoclonal antibodies specific for the amastigote stage of Leishmania pifanoi. I.Characterization of antigens associated with stage- and species-specific determinants' see abstract see page 2407, left column, paragraph 2 see page 2407, right column, last paragraph - page 2408, right column, paragraph 1 see page 2408, right column, paragraph 4 - page 2409, left column, paragraph 1 see page 2409, right column, paragraph 2 - page 2410, right column, paragraph 1 see page 2411, left column, paragraph 3 - right column, paragraph 1 see page 2412, left column, paragraph 2 - right column, paragraph 1 ---	13,18, 24,29,30
A	INFECTION AND IMMUNITY, vol.36, no.1, April 1982, WASHINGTON US pages 430 - 431 K.P. CHANG ET AL. 'Antigenic changes during intracellular differentiation of Leishmania mexicana in cultured macrophages' see abstract see page 431, left column, paragraph 1 - right column, paragraph 1 --- -/--	13



# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/LA 00482

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>MOLECULAR AND CELLULAR BIOLOGY, vol.14, no.5, May 1994 pages 2975 - 2984 HUGUES CHAREST ET AL. 'Developmental gene expression in Leishmania donovani: Differential cloning and analysis of an amastigote-stage-specific gene' see abstract see page 2975, right column, paragraph 3 - page 2976, left column, paragraph 1 see page 2977, right column, paragraph 3 - page 2978, left column, paragraph 1; figure 5 see page 2981, right column, paragraph 2 - page 2982, left column, paragraph 1 -----</p>	1-18

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/CA94/00482

**Box I** Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark : Although claim 29 is directed to a method of treatment of the human/animal body the search has been carried out and based on the alleged effects of the composition.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II** Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant:

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**